


# Plantar Fasciitis



Plantar fasciitis is the most common cause of pain on the bottom of the heel. Approximately 2 million patients are treated for this condition every year.

Plantar fasciitis occurs when the strong band of tissue that supports the arch of your foot becomes irritated and inflamed.

## Anatomy

The plantar fascia is a long, thin ligament that lies directly beneath the skin on the bottom of your foot. It connects the heel to the front of your foot, and supports the arch of your foot.

## Cause

The plantar fascia is designed to absorb the high stresses and strains we place on our feet. But, sometimes, too much pressure damages or tears the tissues. The body's natural response to injury is inflammation, which results in the heel pain and stiffness of plantar fasciitis.

## Risk Factors

In most cases, plantar fasciitis develops without a specific, identifiable reason. There are, however, many factors that can make you more prone to the condition:

- Tighter calf muscles that make it difficult to flex your foot and bring your toes up toward your shin

- Obesity
- Very high arch
- Repetitive impact activity (running/sports)
- New or increased activity

## Heel Spurs

Although many people with plantar fasciitis have heel spurs, spurs are not the cause of plantar fasciitis pain.



Because the spur is not the cause of plantar fasciitis, the pain can be treated without removing the spur.

## Symptoms

The most common symptoms of plantar fasciitis include:

- Pain on the bottom of the foot near the heel
- Pain with the first few steps after getting out of bed in the morning, or after a long period of rest. The pain usually subsides after a few minutes of walking.
- Greater pain after (not during) exercise or activity

## Examination

A comprehensive physical exam is usually enough to diagnose plantar fasciitis.

**OAM+** Orthopaedic Associates of Muskegon

**West Michigan Spine Center**

**Grand Haven Bone & Joint**

1400 Mercy Drive, Ste 100  
Muskegon, MI 49444  
231-733-1326

1445 Sheldon Rd, Suite G1  
Grand Haven MI 49417  
616-296-9100

[www.oamkg.com](http://www.oamkg.com)  
[www.wmspinecenter.com](http://www.wmspinecenter.com)

## X-rays

X-rays provide clear images of bones. They are useful in ruling out other causes of heel pain, such as fractures or arthritis. Heel spurs can be seen on an x-ray. MRI is rarely needed and only ordered if other causes of pain are of concern.

## Nonsurgical Treatment

More than 90% of patients with plantar fasciitis will improve within 10 months of starting simple treatment methods.

**Rest:** Decreasing or even stopping the activities that make the pain worse is the first step in reducing the pain. You may need to stop athletic activities where your feet pound on hard surfaces (for example, running or step aerobics).

**Ice:** Rolling your foot over a cold water bottle or ice for 20 minutes is effective. This can be done 3-4 times a day.

**Nonsteroidal anti-inflammatory medication:** Drugs such as ibuprofen or naproxen reduce pain and inflammation. Using the medication for more than 1 month should be reviewed with your primary care doctor.

**Exercise:** Plantar fasciitis is aggravated by tight muscles in your feet and calves. Stretching your calves and plantar fascia is the most effective way to relieve the pain that comes with this condition.

- **Calf stretch:** Lean forward against a wall with one knee straight and the heel on the ground. Place the other leg in front, with the knee bent. To stretch the calf muscles and the heel cord, push your hips toward the wall in a controlled fashion. Hold the position for 10 seconds and relax. Repeat this exercise 20 times for each foot. A strong pull in the calf should be felt during the stretch.
- **Plantar fascia stretch:** This stretch is performed in the seated position. Cross your affected foot over the knee of your other leg. Grasp the toes of your painful foot and slowly pull them toward you in a controlled fashion. If it is difficult to reach your foot, wrap a towel around your big toe to help pull your toes toward you. Place your other hand along the plantar fascia. The fascia should feel like a tight band along the bottom of your foot when stretched. Hold the stretch for 10 seconds. Repeat it 20 times for each foot. This exercise is best done in the morning before standing or walking.



injected into the plantar fascia to reduce inflammation and pain. However, multiple steroid injections can cause the plantar fascia to rupture (tear), which can lead to a flat foot and chronic pain.

**Supportive shoes and orthotics:** Shoes with thick soles and extra cushioning can reduce pain with standing and walking. A cushioned shoe or insert reduces this tension and the microtrauma that occurs with every step. Soft silicone heel pads are inexpensive and work by elevating and cushioning your heel. Pre-made or custom orthotics (shoe inserts) are also helpful.



Soft heel pads can provide extra support.

**Night splints:** Most people sleep with their feet pointed down. This relaxes the plantar fascia and is one of the reasons for morning heel pain. A night splint stretches the plantar fascia while you sleep. Although it can be difficult to sleep with, a night splint is very effective and does not have to be used once the pain is gone.

**Physical therapy:** Your doctor may suggest that you work with a physical therapist on an exercise program that focuses on stretching your calf muscles and plantar fascia. In addition to exercises like the ones mentioned above, a physical therapy program may involve specialized ice treatments, massage, and medication to decrease inflammation around the plantar fascia.

## Surgical Treatment

Surgery is considered only after 12 months of aggressive nonsurgical treatment and is very rarely indicated.

**Gastrocnemius recession:** This is a surgical lengthening of the calf (gastrocnemius) muscles. Because tight calf muscles place increased stress on the plantar fascia, this procedure is useful for patients who still have difficulty flexing their feet, despite a year of calf stretches.

**Plantar fascia release:** If you have a normal range of ankle motion and continued heel pain, your doctor may recommend a partial release procedure. During surgery, the plantar fascia ligament is partially cut to relieve tension in the tissue. If you have a large bone spur, it will be removed, as well.

**Recovery:** Most patients have good results from surgery. However, because surgery can result in chronic pain and dissatisfaction, it is recommended only after all nonsurgical measures have been exhausted.

*Adapted from American Academy of Orthopaedic Surgeons. For more information, see [orthoinfo.aaos.org](http://orthoinfo.aaos.org)*

**Cortisone injections:** Cortisone, a type of steroid, is a powerful anti-inflammatory medication. It can be